

Ground state properties of dissipative two state system coupled to phonons

T. P. Pareek¹ and A. M. Jayannavar²

Institute of Physics, Bhubaneswar-751005, India

February 1, 2008

ABSTRACT

We present a systematic analysis of ground state properties of a tunneling system coupled to an Ohmic bath. The nonadiabatic effects induced by coupling with tunneling particle , on the ground state properties of phonons are taken into account. For this we have considered both displacement and deformation of phonons via a variational treatment. The symmetry breaking transition , or the delocalization-localization transition of tunneling particle

¹e-mail: pareek@iopb.ernet.in

²e-mail: Jayan@iopb.ernet.in

is presented and is compared with earlier known results. We also show that in the critical regime a gap will be opened near the zero momentum in the spectrum of the phonons. The modified phonon spectrum along with the phonon displacement field correlation functions have been calculated. It is shown that in the critical regime the tunneling system stabilizes the lattice against long wavelength fluctuations.

PACS NO: 73.40.Gk , 74.50.+r , 03.65 - ω

1. Introduction

The statics and dynamics of two state tunneling system coupled to an environment is of fundamental interest both in physics and chemistry [1]. This rather simple system has been used to study the motion of impurities (atoms or groups of atoms) in solids and glasses which play an important role in determining low temperature properties [2,3], defect motion in alkali halides [4], charged particle motion in metals, magnetic impurities in metals [5,6] and molecular transition in liquids [7] etc. In all examples mentioned above tunneling is a many-body problem in the sense that the considered particle couples to a large number of degrees of freedom arising from medium or environment. In recent years there has been a growing interest in the influence of dissipation on the behaviour of macroscopic quantum variables and quantum tunneling out of metastable state. In particular, the trapped magnetic flux threading a superconducting ring interrupted by weakly coupled Josephson junction is expected to show macroscopic quantum coherence. In this case the problem of tunneling of quantized magnetic flux reduces to quantum motion of a particle in a double well potential [1]. The existence of chiral molecules by a superselection rule which originates from the ever

present coupling of the molecule to the radiation field can be understood in the frame work of two state system [8]. Here the two states are nothing but the left and right handed configurations of molecule. Recently the two state system formulation has been applied to an understanding of c-axis resistivity in high- T_c layered oxides and transverse resistivity in quasi one-dimensional compounds [9,10] . Electrical transport along c-axis in layered oxides is pictured as a coherent interplanar tunneling between neighbouring layers blocked by repeated intraplanar incoherent(inelastic) scattering. Here two states correspond to two nearest neighbour layers. The ground state properties of two level system are also related to anisotropic Kondo problem [1] and one dimensional inverse square Ising model [11,12] . In particular the delocalization-localization transition in two state system is related to the existence of ferromagnetic phase transition in the one-dimensional inverse square Ising model and several rigorous results for this problem are known.

In all these systems it is now well established that the environmental coupling to tunneling system, makes a tunneling motion less frequent and consequently reduces the effective tunneling matrix element, sometimes termed as Debye-Waller factor. This reduction stems from orthogonal properties of many particle wavefunctions with different local potentials and is known as

Anderson orthogonality theorem [13] . In our study we examine the ground state properties of two state system linearly coupled to a phonon bath which provides Ohmic dissipation. This system can be described by a spin-boson Hamiltonian.

$$H = -\Delta_o \sigma_x + \sum_k \omega_k a_k^\dagger a_k + \sum_k g_k (a_k + a_{-k}^\dagger) \sigma_z \quad , \quad (1)$$

where Δ_o represents bare tunneling matrix element, σ_x, σ_y are Pauli matrices. The bath is described by a set of harmonic oscillators and with coupling constant g_k and $a_k, a_k^\dagger$ are the annihilation and creation operators representing phonon modes. We have taken coupling constant g_k of the form $g_k = g_o |k|^{1/2}$, which represents coupling of a tunneling system to a one-dimensional phonons. The frequency dispersion is taken to be $\omega_k = |k|$ (acoustic phonons) , where k is the quasimomentum wavevector of phonons. The Hamiltonian(1) can also be written in terms of creation ($c_i^\dagger$) and annihilation (c_i) operators of the tunneling particle, namely

$$H = -\Delta_o (c_1^\dagger c_2 + c_2^\dagger c_1) + \sum_k \omega_k a_k^\dagger a_k + \sum_k g_k (c_1^\dagger c_1 - c_2^\dagger c_2) (a_k + a_{-k}^\dagger) \quad , \quad (2)$$

where 1 and 2 represent two equivalent positions of tunneling particle. Since we are dealing with a single particle it does not matter whether it is a boson or a fermion. The Hamiltonian for two state system is also a good representation

for statics and dynamics of a quantum particle moving in symmetric double well potential ,i.e, in the limit of low temperatures, $k_B T$ much smaller than $\hbar\omega_o$ (ω_o the frequency of small amplitude oscillations in the wells) and V_o the barrier height between the wells [1]. The tunneling frequency Δ_o is of the order of $[\sqrt{\omega_o V_o/\hbar}]exp(-2V_o/\hbar\omega_o)$, in the WKB limit. The Ohmic dissipative bath corresponds to a special case [1] where the spectral density $S(\omega)$ is given by

$$S(\omega) = \sum_k g_k^2 \delta(\omega - \omega_k) = \frac{1}{2} \alpha \omega \quad , \quad (3)$$

where α is the dimensionless dissipation parameter.

The above Hamiltonian exhibits a rich phenomena for a tunneling system ranging from damped oscillations(coherence) up to localization [1] . It has been shown that for the weak coupling the particle tunnels back and forth between two states and quickly loses the memory of the initial quantum state. As the coupling exceeds the critical strength , spontaneous symmetry breaking occurs , leading to localization where the effective tunneling matrix is identically zero [1,14,15] . This symmetry breaking transition occurs only in the limit of zero temperature. At finite temperature and for larger coupling

strength the tunneling particle hops incoherently from one state to another. The zero temperature localization phenomenon is associated with the infrared divergence induced by low energy phonons ,i.e, the phonon field keeps the particle localized at the expense of generating an infinite number of low-energy phonons. The low frequency phonons play a dominant role in the statics as well as in the dynamics of the two-state system. The conventional weak coupling picture has a rather limited range of validity and the adiabatic approximation over estimates the occurrence of infrared divergence.

Using the renormalization-group procedure the delocalization to localization transition has been studied [1]. When the bare tunneling matrix element Δ_o is small compared to the upper phonon cutoff frequency ω_c , the effective tunneling matrix element is given by $\Delta_{eff} = \Delta_o (\frac{2e\Delta_o}{\omega_c})^{\frac{\alpha}{(1-\alpha)}}$. At $\alpha = 1$ there is a transition from delocalized to localized ($\Delta_{eff} = 0$) state. The renormalization-group results have been obtained mainly based on the assumption of a dilute instanton gas or dilute non-interacting blip approximation [1] . In the later studies on dissipative quantum tunneling , it has been shown that this approximation breaks down due to the condensation of instanton gas [16] . It has also been pointed out that the non-interacting blip approximation leads to a complete neglect of reaction field which becomes

important for the long time dynamics in the larger coupling domain [17]. Silbey and Harris [18], and Tanaka and Sakurai [19] have independently succeeded in rederiving the renormalization-group results by a simple variational procedure which amounts to a kind of mean field theory. They have taken the displacement of phonons in a ground state via variational parameters. It should be noted that when a particle tunnels from one state to another it has two distinct effects on the phonon wave function. One is that of a displacement and the other is that of a deformation. High-frequency phonons ($\omega > \Delta_o$) follow the tunneling particle adiabatically. Hence the displaced phonons gives the correct result in this domain. However, for low-frequency modes and in the strong coupling regime such an approximation is not sufficient. The low frequency phonons in general are not always in equilibrium with the particle motion, and hence the tunneling particle not only moves in an adiabatic potential, but also gets influenced by a retarded potential arising due to the phonon bath [4,21]. The motion of the particle at time t' disturbs the phonon state, which in turn acts on the motion of the particle at later time t . For the frequencies $\omega < \Delta_o$, these non-adiabatic effects dominate, which in turn apart from displacement produce strong deformation in the ground state phonon wave function. Chen and coworkers have

treated these effects via a variational treatment by introducing a squeezing parameter for the phonon states [20,22-24]. Within this treatment they have analysed the symmetry breaking transition and have also shown that the ground state properties of the tunneling particle not only depends on the spectral density , but also on the explicit form of the coupling strength on frequency of phonon modes [20]. This non-universal behaviour is at variance with the earlier results based on the renormalization-group and the obtained phase diagram is in no simple way related to the known results [1]. In Ref[25] , the phase diagram for the delocalization-localization transition has been analysed through a variational treatment which incorporates both the displacement and the squeezing of wave function via two independent variational parameters for each phonon mode.

Most of the theoretical treatments to date are concerned with the effect of phonon bath on the tunneling motion. Very few treatments exist which directly address the question of effect of tunneling particle on the phonon bath [20,26,27]. It has been shown that [20] two state coupled system at zero temperature can be mapped on to the modified quantum sine-Gordon model , which describes the effective non-linear interaction between the low frequency phonons when the tunneling system is in its ground state. It is also

of interest to know that the ground state properties of $(1 + 1)$ dimensional quantum sine-Gordon model can be mapped on to several two dimensional statistical mechanics models ,e.g., two dimensional Coulomb gas etc [27]. In the critical regime , where the particle is still in the delocalized state , a gap in the elementary excitation of the low frequency phonons is opened near zero momentum , and the corresponding ground state wave function is found to be of a pairing quasi-particle state , analogous to the BCS super conducting state [26,27].

In our present work we have carried out a systematic analysis of ground state properties of the tunneling particle as well as of the phonons via a variational treatment which correctly treats both the displacement and deformation of each phonon wave function in its ground state . As stated earlier we have restricted our treatment to a two state system , coupled to bosonic environment, with a Ohmic spectral density. Results do depend on specific form of dependence of coupling constant on the frequency of phonon mode, in our case $g_k = g_o\sqrt{\omega_k}$, this coupling is common to electron-phonon coupling in solids. The reduction in the effective tunneling matrix element of the particle as a function of dimensionless dissipation parameter α is calculated and is compared with earlier known results. The modified phase diagram for the

symmetry breaking transtion is presented. We have also shown that in the critical regime and in the delocalized phase the gap will be opened near zero momentum in the spectrum of the low frequency phonons. The gap vanishes throughout in the localized domain. An analytical expression for the phonon frequency gap as a function of effective tunneling matrix element and dissipation parameter α is obtained. We have also obtained an information about the modified phonon spectrum and using this , the phonon displacement field spatial correlation function has been calculated. In the delocalized phase the phonon field correlation functions at large distances (in the asymptotic domain) falls of exponentially. This implies that in the presence of a tunneling particle the lattice is stable under the long wave length fluctuations , so long as effective tunneling matrix element for a particle is non-zero.

2. Ground state variational wave function

In the following we first motivate our ground state variational wave function of the system described by the Hamiltonian given in eqn (1) or (2) . Ground state wave function for this system can not be obtained analytically. Only special cases can be solved exactly. The simple cases being (i) $\Delta_o = 0$, and (ii) the coupling constant $g_k = 0$. In the first case where $\Delta_o = 0$, the

system is just an oscillator and there is no tunneling motion between the two degenerate states. However, the oscillators are displaced in one direction when the tunneling system is in one of the two levels and are displaced in the opposite direction when the tunneling system is in the other level. Correspondingly the total system in its ground state can be represented by a doubly degenerate localized state

$$\phi_i = \prod_k \exp(-\frac{1}{2} \frac{g_k^2}{\omega_k^2}) \exp\{(-1)^i \frac{g_k}{\omega_k} b_k^\dagger\} |0\rangle \quad . \quad (4)$$

Here $|0\rangle$ represents the vacuum state and i being the position of the tunneling state. For an Ohmic dissipative bath the overlap integral between the above two states tends to zero, if there is no lower cutoff frequency, i.e., the two ground states are orthogonal to each other. In the other cases where $g_k \neq 0$, i.e., tunneling particle and phonon modes decouple, the eigenstates of the tunneling particle are given by the symmetric and antisymmetric combination of two level states, the symmetric combination being the ground state. These two situations indicate that for a finite value of g_k and Δ_o the system exhibits the competition between the localization resulting from the phonon modes and the delocalization due to the tunneling. The particle tunnels from one

state to another , carrying a dressed phonon cloud. In view of these facts , we immediately write down a ground state variational wave function as

$$\psi = \frac{1}{\sqrt{2}} \left\{ \prod_k \exp\left[-\frac{1}{2} |u_k|^2\right] \exp[-u_k a_k^\dagger] \exp[-\gamma_k(a_k a_{-k} - a_k^\dagger a_{-k}^\dagger)] C_1^\dagger + \prod_k \exp\left[-\frac{1}{2} |u_k|^2\right] \exp[u_k a_k^\dagger] \exp[-\gamma_k(a_k a_{-k} - a_k^\dagger a_{-k}^\dagger)] C_2^\dagger \right\} |0\rangle \quad , \quad (5)$$

where u_k and γ_k are both variational parameters for each phonon mode. We call our variational wave function (eqn.[5]) as VWF(3) . When the parameter γ_k takes a finite non zero value we say that the corresponding phonon state (or two phonon state) is squeezed [28]. The squeezed state represents a non-classical minimum uncertainty state and are eigenstates of a linear combination of the annihilation and creation operators. It differs from a minimum uncertainty coherent state in that it has uncertainty less than that of a coherent state in one of the quadrature phase given by $\exp(-\gamma_k)$ and in another $\exp(\gamma_k)$, which means that one of the quadrature phase is squeezed at the expense of other. In real space representation the displaced squeezed state is given as $\phi(x) = [\pi S^2]^{-1/4} \exp[-(x-x_0)^2/2S^2]$, where x_0 is the mean displacement and $S = \exp(\gamma_k)$. For this state the uncertainty , Δx in spatial coordinate equals to $\exp(\gamma_k)$ and Δp in the momentum equals $\exp(-\gamma_k)$. If

γ_k takes a positive value the wave function spreads over a larger distance than the ground state wave function of the harmonic oscillator (coherent state). Squeezing parameter takes care of the deformation associated with the low frequency phonon wave function as discussed earlier. In VWF(3) if we substitute $\gamma_k = 0$, we get the variational wave function proposed independently by Tanaka and Sakurai [19] and Silbey and Harris [18], we call this VWF(1). In the VWF(1), if we substitute $u_k = 0$, it becomes an exact ground state of the unperturbed tunneling particle. On the other hand if we put $u_k = -g_k/\omega_k$, the wave function describes a linear combination of two degenerate ground states given in eqn(4). The VWF(1) does not take into account the deformation of low frequency phonons. In VWF(3) if we take variational parameter u_k to be constant and equal to $(-g_k/\omega_k)$, we get the variational wave function proposed by Chen et.al [20,22-24]. We call this as a VWF(2). The VWF(2) takes in to account the deformation of low frequency phonon via squeezing parameter γ_k . However, it does not take properly in to account the displacement of phonons. All the phonons are assumed to be displaced by an amount $|u_k| = g_k/\omega_k$, which is true only in the adiabatic limit or for phonon frequency $\omega > \Delta_o$. In the VWF(3) both u_k and γ_k being the variational parameter, naturally it takes care of non adia-

batic effects in both the displacement and the deformation of low frequency phonons properly.

Now using VWF(3) one can obtain the expectation value of the total ground state energy (E_g) . Using the well known properties [20,22-30] of squeezed states and bose operators we get,

$$E_g(u_k, \gamma_k) = -\Delta_o \left\{ \exp \sum_k [(-2u_k^2) \exp(-4\gamma_k)] \right\} + \sum_k \omega_k [u_k^2 + (\sinh 2\gamma_k)^2] + \sum_k 2g_k u_k \quad . \quad (6)$$

The variational parameters u_k and γ_k are determined by minimising E_g with respect to u_k and γ_k , i.e. , we set $\frac{\partial E_g}{\partial u_k} = 0$ and $\frac{\partial E_g}{\partial \gamma_k} = 0$. From this we obtain

$$u_k = [-g_k / (2\Delta_o Z \exp(-4\gamma_k) + \omega_k)] \quad , \quad (7)$$

$$\gamma_k = (1/8) \ln \left[1 + 8\Delta_o Z u_k^2 / \omega_k \right] \quad . \quad (8)$$

The factor multiplying Δ_o in the eqn. (6) is identified as a tunneling reduction factor Z (or Debye-Waller factor), and is given by

$$Z = \exp \left\{ \sum_k [-2u_k^2 \exp(-4\gamma_k)] \right\} \quad . \quad (9)$$

The effective matrix element Δ_{eff} is given by $\Delta_{eff} = \Delta_o Z$. The reduction factor Z can be determined by using eqns.(8) and (9) self-consistently. One

can notice from eqns.(7) and (8) that u_k and γ_k tends to $(-g_k/\omega_k)$ and zero respectively in the high frequency limit as discussed earlier. This is consistent with an adiabatic treatment in this frequency domain , i.e. , high frequency phonons with ω greater than the Δ_{eff} follow the tunneling particle adiabatically.

3. Effective phonon Hamiltonian

We will now derive an effective Hamiltonian for phonon bath at zero temperature consistent with the VWF(3). To this end we first perform a canonical transformation

$$U_1 = \exp \left[\sigma_z \sum_k u_k (a_k - a_{-k}^\dagger) \right] \quad , \quad (10)$$

on Hamiltonian (1). we get

$$\begin{aligned} H_1 = U_1^\dagger H U_1 = & \sum_k \omega_k a_k^\dagger a_k - \Delta_o \sigma_x \text{Cosh} \left[\sum_k 2u_k (a_k - a_{-k}^\dagger) \right] \\ & + i\Delta_o \sigma_y \text{Sinh} \left[\sum_k 2u_k (a_k - a_{-k}^\dagger) \right] \\ & + \sum_k (g_k + u_k \omega_k) \sigma_z + 2 \sum_k (u_k^2 \omega_k + g_k u_k) \quad . \end{aligned} \quad (11)$$

From the Hamiltonian in eqn.(11) we see that there are two distinct types of influence on the phonon bath due to coupling with two-state system. The

first is the diagonal term containing σ_x , giving static influence due to two level system being its ground state ($\sigma_x = +1$). The nondiagonal interaction terms containing σ_y and σ_z represents the dynamic influence due to the transition in two state system between its ground state and excited state on the phonon system. These nondiagonal terms are important only at finite temperatures. Hence at zero temperature we neglect these terms along with the constant factor appearing in eqn.(11), thus we have

$$H_{eff} = \sum_k \omega_k a_k^\dagger a_k - \Delta_o \cosh \left[\sum_k 2u_k (a_k - a_{-k}^\dagger) \right] \quad . \quad (12)$$

Equation.(12) represents an effective phonon Hamiltonian, having a nonlinear interactions between the phonons. Here nonlinear interactions are not only between the phonon in the same mode, but also in the different modes. By appropriately defining the field operators one can readily map the H_{eff} on to modified quantum sine-Gordon model (see ref[20]) in 1+1 space-time dimension. It is well known that the quantum sine-Gordon model exhibits a Coleman phase transition as a function of the coupling constant. In Ref[26,27] critical conditions for this phase transition have been derived. It is also shown that, in the critical regime the gap will be opened in the excitation spectrum near zero momentum. The Ref[27], also elaborates on the connection with

Kosterlitz-Thouless classical theory of two dimensional Coulomb gas. In these references [20,26,27] static properties of two state systems have been studied , by mapping to sine-Gordon model. The results obtained are in spirit of VWF(2). However , our results differ in that we closely follow the spirit of VWF(3) , which itself is a better ground state variational wave function. For the effective phonon system (eqn.[12]) , we apply the second unitary transformation

$$U_2 = \exp \left[\sum_k \gamma_k (a_k a_{-k} - a_k^\dagger a_{-k}^\dagger) \right] . \quad (13)$$

This transforms the effective Hamiltonian (12) into

$$\begin{aligned} H_{eff} = & \sum_k \omega_k [b_k^\dagger b_k \cosh 4\gamma_k + (\sinh 2\gamma_k)^2 \\ & + (1/2)(b_k^\dagger b_{-k}^\dagger + b_k b_{-k})(\sinh 4\gamma_k)] \\ & - \Delta_o \cosh \left[\sum_k 2u_k (b_k - b_{-k}^\dagger) \exp(-2\gamma_k) \right] , \end{aligned} \quad (14)$$

where b_k and $b_k^\dagger$ are

$$b_k = U_2^\dagger a_k U_2 = a_k \cosh 2\gamma_k - a_{-k}^\dagger \sinh 2\gamma_k , \quad (15)$$

$$b_k^\dagger = U_2^\dagger a_k^\dagger U_2 = a_k^\dagger \cosh 2\gamma_k - a_{-k} \sinh 2\gamma_k . \quad (16)$$

The transformation (13) some times referred to as a Bogolibov transformation, basically it rescales the phonon normal-mode coordinate and momenta. We

can expand the hyperbolic function in eqn.(14) by normal ordering phonon operators. We choose γ_k such that it minimises the ground state energy as given by eqn.(8). With this choice of γ_k we get , after the straight-forward algebra (see also ref.[27])

$$H_{eff} = \sum_k \omega_k \exp(4\gamma_k) b_k^\dagger b_k + E_o - \Delta_o Z [(i\phi_+)^4/240 - (i\phi_+)^3(i\phi_-)/6 +] \quad , \quad (17)$$

where ϕ_+ and ϕ_- are

$$\phi_+ = \sum_k u_k \exp(2\gamma_k) b_{-k}^\dagger \quad , \quad (18)$$

$$\phi_- = \sum_k u_k \exp(-2\gamma_k) b_k \quad . \quad (19)$$

The ground states of (17) are displaced squeezed states in the original phonon bases.

$$\phi = \exp\{\sum_k \gamma_k (a_k a_{-k} - a_k^\dagger a_{-k}^\dagger)\} \exp\{-\sigma_z \sum_k u_k (a_{-k}^\dagger - a_k)\} | 0 \rangle \quad . \quad (20)$$

We see that Hamiltonian (17) corresponds to a free field model if the tunneling reduction factor Z approaches zero. The effective tunneling matrix element approaching zero implies the symmetry breaking transition for a tunneling particle in that the particle goes from the delocalized state to a localized state .The effective tunneling matrix element $\Delta_{eff} \rightarrow 0$ also implies

the vanishing of nonlinear interaction terms in Hamiltonian (17) as well as in the Hamiltonian (12) . In eqn. (17) we identify $\Omega(k) = \omega_k \exp(4\gamma_k)$ as the new excitation spectrum of the phonon quasiparticles with their ground state wave function given by eqn.(20) . It is clear that in the critical regime where Z is small ($Z \ll 1$) , the higher order terms in eqn.(17) are small and correspondingly $\Omega(k)$ is well defined excitation spectrum of a modified phonon bath in the presence of a tunneling particle. In this limit we later show that the gap will be opened in the vicinity of the zero momentum in the excitation spectrum of the phonons and this gap will disappear as $Z \rightarrow 0$. The opening up of the gap in the vicinity of the zero momentum naturally takes care of infrared divergence problem arising in a effective tunneling matrix element without invoking infrared cutoff prescription. With a modified excitation spectrum near the critical regime , we will also calculate the phonon displacement field correlation functions.

4. Numerical procedure

We will first outline the numerical technique used to investigate the behaviour of the reduction factor from the expression for Z (eqn[9]) . For the

case of Ohmic dissipation this can be written as

$$\ln Z = -\alpha \int_0^{\omega_c} d\omega \omega \exp(-4\gamma_k) / (2\Delta_o Z \exp(-4\gamma_k) + \omega)^2 \quad , \quad (21)$$

where $\alpha = g_o^2/\pi$ is a dimensionless dissipation parameter, which characterises the coupling between the particle and the surrounding medium and ω_c is the upper phonon frequency cut-off. Throughout we have taken $\omega_k = |k|$. To evaluate Z we have to use u_k and γ_k as given in eqns.(7) and (8) respectively. The expression for γ_k contains u_k and vice-versa. Substituting these values for u_k and γ_k in eqn.(21) leads to an infinite (continued fraction) expression. In ref [25] , this self-consistent equation has been solved by iterative procedure where one terminates the infinite heirarchy to a finite one. This procedure is a approximate one. We have developed an exact numerical procedure without resorting to any approximations. To this end we define a new quantity $\Omega(k) = \omega_k \exp(4\gamma_k)$ with this definition eqn.(21) reduces to

$$\ln Z = -\alpha \int_0^1 d\omega \Omega(k) / (2\Delta_o Z + \Omega(k))^2 \quad . \quad (22)$$

Using eqns (7) and (8) one can readily show that $\Omega(k)$ obeys a forth order polynomial equation , namely

$$\Omega^4 + 4\Delta_0 Z \Omega^3 + (4\Delta_0^2 Z^2 - 8\Delta_0 Z g_k^2 / \omega_k - \omega_k^2) \Omega^2 + (4\omega_k^2 \Delta_0 Z) \Omega - 4\omega_k^2 \Delta_0^2 Z^2 = 0 \quad . \quad (23)$$

Here $g_k = g_o | \omega_k |^{1/2}$. In eqns.(22) and (23) we have rescaled the energy with respect to ω_c , for example $\Delta_o = \Delta_o / \omega_c$, $\Omega(k) = \Omega(k) / \omega_c$ etc. For a given value of $\alpha(= g_k^2 / \pi)$, and Z this equation gives four solutions. It turns out that out of these four solutions two are complex and one is positive and another is negative. We take only the physical solution, i.e., positive one as it represents the energy of the modified phonon modes. In some parameter range where Δ_o / ω_c is large, four solution to Ω of eqn.(23) are all real. It turns out that three of them are negative, hence we again take a positive solution. We first take a fixed value of α , and Δ_o and for a value of Z lying in an interval $1 < Z < 0$ we calculate $\Omega(k)$ for all ω_k . Substituting this $\Omega(k)$ for all ω_k in the expression (22) we check for self-consistency. We start initially with $Z=1$ and go on changing Z in a small step till we reach the self-consistent solution.

5. Results and discussion

In fig.(1) we have plotted the reduction factor Z as a function of dimensionless parameter α for a fixed value of $\Delta_o/\omega_c = 0.1$. For comparison we have also plotted the earlier results obtained from VWF(1) (....) and VWF(2) (- - -). In all these cases the reduction factor decreases as expected as a function of α and at a particular value α_c there is a symmetry breaking transition from delocalized state to localized state, at which Z becomes identically zero. For $\Delta_o/\omega_c = 0.1$ the VWF(1) gives α_c around 1, whereas VWF(2) gives α_c around 2. In our case we obtain $\alpha_c = 1.85$. We have identified α_c such that at this value Z becomes less than 10^{-7} . In parameter range of α between zero and 0.5 the results obtained from VWF(1) and VWF(3) are almost identical. As we approach α_c the results deviate. This deviation should be identified as a result of inclusion of fluctuations over the mean field results obtained earlier [1]. It is also of interest to note that the parameter range $0 < \alpha < 0.5$ in the limit $\Delta_o/\omega_c \ll 1$ is of most interest for the quantum coherence problem [14]. In this regime at temperature $T=0$, the tunneling system exhibits damped oscillations (coherence) with frequency Δ_{eff} . The symmetry breaking transition obtained from VWF(3) is continuous in the

sense that the reduction factor Z approaches zero continuously as α tends to the critical value α_c . Even for larger values of Δ_o/ω_c we have observed that the results obtained for the reduction factor Z from VWF(1) and VWF(3) are identical in the range of small dissipation parameter $\alpha \ll 1$. In the same parameter range of α the results obtained from VWF(2) and VWF(3) differ considerably. Moreover, it should be noted that the results for Z obtained from VWF(1) in the limit $\alpha \ll 1$ are consistent with the known perturbation results [17].

In fig(2) we have plotted the phase boundary between the delocalized ($Z \neq 0$) and localized ($Z = 0$) phase for various values of Δ_o/ω_c . In the limit of $\Delta_o/\omega_c \rightarrow 0$ the phase boundary is located at $\alpha_c = 1$. The result based on instanton calculation or path integral calculation also predict the phase boundary to be at $\alpha_c = 1$ as $\Delta_o/\omega_c \rightarrow 0$. It also predicts the phase boundary to be at $\alpha_c = 1$ till Δ_o/ω_c reaches a value of $1/2e = 0.184$. In our case the phase boundary shifts to the right side (increase in α_c) continuously as we vary Δ_o/ω_c . The VWF(2) predicts the phase boundary starting from $\alpha_c = 2$ in the limit $\Delta_o/\omega_c \rightarrow 0$ to a higher values of α_c as Δ_o/ω_c increases. In a sense the phase boundary of the delocalization to localization transition obtained from VWF(1) and VWF(2) are not related. We can say with a

certain confidence that our results clearly shows the improvement over the earlier results obtained by various treatment using path integral formulation [1]. In effect our results amount to taking care of flucutations (via a squeezing parameter) in an otherwise mean field results obtained earlier.

From eqn.(17) we have identified $\Omega(k) = |k| \exp(4\gamma_k)$ as excitation spectrum of the modified phonon bath under the coupling with a tunneling particle. This excitation spectrum aquires a well defined meaning only when the Δ_{eff} is small , in this regime the higher order correction terms in eqn.(17) becomes quite small and give a small correction to the excitation spectrum via perturbation. In fig.(3) we have plotted the excitation spectrum $\Omega(k)$ versus wave vector k , for $\Delta_o/\omega_c = 0.1$ and $\alpha = (.9599)$ for which $Z=.1005$. We have calculated the excitation spectrum with the help of eqn.(23) for a given value of α and corresponding value of Z which has been calculated by self consistent procedure as mentioned earlier. The eqn.(23) for $\Omega(k)$ gives four solutions and out of these there is only one physically acceptable solution which gives a positive value of $\Omega(k)$. We see from the fig.(3) that the gap opening in the vicinity of zero momentum. For comparision we have also plotted the excitation spectrum $\omega_k = |k|$ of the unperturbed free phonon modes (in the absence of coupling to a tunneling system). It is clear from

the figure that the low frequency phonon modes are affected due to coupling as compared to high frequency modes as expected. In fig.(4) we have plotted the zero momentum excitation gap $\Omega(0)$ versus the dimensionless dissipative constant α , for a fixed value of $\Delta_o/\omega_c = 0.1$. The gap has initially a value zero up to some α , then it increases and after exhibiting a maxima the gap vanishes identically at the critical value of α_c . In the regime where the gap $\Omega(0)$ is finite, it is given by $\Omega(0) = -2\Delta_o Z + 2\sqrt{(2\Delta_o Z \alpha \pi)}$. Even though in fig.(4) we have plotted $\Omega(0)$ versus α for the full range of α where tunneling system undergoes delocalization to localization transition, the gap $\Omega(0)$ is meaningful quantity only in the critical regime, i.e., α close to α_c ($Z \ll 1$). If one would have carried out the calculations in the spirit of VWF(2), i.e., by setting the parameter $u_k = (-g_k/\omega_k)$ in the canonical transformation given in eqn.(10), we can readily obtain [26,27] an analytical expression for the excitation spectrum of the modified phonon bath. This is given by $\Omega(k) = |k| [1 + (4\pi\alpha\Delta_o Z/k^2)]^{1/2}$, with a gap in the vicinity of zero momentum $\Omega(0) = \sqrt{4\pi\alpha\Delta_o Z}$. The expression for $\Omega(k)$ depends explicitly on α, Z, Δ_o , and k . As the results derived from VWF(2) and VWF(3) for Z and consequently the symmetry breaking transition as a function of α differ qualitatively, we have not made the comparison between $\Omega(k)$ obtained in

the spirit of VWF(2) and VWF(3).

Once we have the full information about the spectrum one can easily compute the phonon displacement field spatial correlation function $C(x)$. This correlation function is defined as [27]

$$C(x) = \frac{1}{2} \int_0^{k_c} \frac{dk}{2\pi} \frac{\cos(kx)}{\Omega(k)}. \quad (24)$$

where k_c is the upper phonon frequency cutoff $k_c = \omega_c$. In fig.(5) we have plotted $C(x)$ versus x ($=xk_c$) for $\Delta_o/\omega_c = 0.1$, $\alpha = (.9599)$ and $Z = (.1005)$. The correlation function initially follows a logarithmic behaviour between the two space points with the increasing of their space distance x . At the large distances (or asymptotically) $C(x)$ decays exponentially, after showing an oscillatory behaviour in the intermediate regime. The characteristic decay length ξ (or the correlation length) is inversely proportional to $\Omega(0)$ and hence ξ becomes large in the critical regime and diverges at α_c , the transition point of delocalization to localization transition. We have also observed on the expanded scale that the domain of $x \ll \xi$, the correlation function decays logarithmically. It is important to note that in the absence of coupling of tunneling particle to lattice phonons, the phonon field correlation function

diverges logarithmically. This indicates the instability of lattice against long wave length fluctuations. However ,by coupling lattice to a tunneling system, the lattice acquires stability against the long wave length fluctuations so long as tunneling system is in a delocalized state (i.e , when the excitation spectrum has a gap in the vicinity of a zero momentum). This fact manifests itself in exponential decay of $C(x)$ in the asymptotic regime . Thus a single impurity (or a tunneling system) can stabilize a one dimensional lattice. The correlation length ξ scales as $\xi \sim Z^{-1/2}$ and it diverges as we approach α_c from below.

In conclusion we have studied the ground state properties of a coupled spin-boson system i.e., a tunneling system coupled to an Ohmic dissipative interaction arising due to a linear interaction of tunneling system to a one-dimensional set of harmonic phonons with an appropriate spectral density. We have presented a modified phase diagram for symmetry breaking transition. The results for the behaviour of reduction factor on the dissipation parameter are given and compared with known results. We have also shown that the effective phonon system shows a gap in the excitation spectrum in the critical regime and phonon displacement field correlation function decays exponentially in the same regime , this stabilizes lattice against the long wave

length fluctuations. Throughout our analysis we have restricted to a special form of coupling constant , namely g_k is proportional to $|k|^{1/2}$. It is known that the different forms for coupling constant lead to qualitatively different results for symmetry breaking transition , in a sense there is no universality in a dissipative two state system [21]. The zero temperature properties depend not only on the phonon spectral density , but also on the explicit dependence of coupling strength on the phonon frequency. We are at present exploring the behaviour of effective phonon spectrum for different dependence of coupling constant and the result will be presented elsewhere.

References

- [1] A. J. Leggett, S. Chakravarty, A. T. Dorsey, Matthew P. A. Fisher and Anupam Garg Rev. Mod. Phys. **59**, 1 (1987).
- [2] P. W. Anderson Philos. Mag. **25**, 1 (1972).
- [3] W. A. Phillips. J. Low Temp. Phys. **7**, 351 (1972).
- [4] J. P. Sethna. Phys. Rev. B **24**, 698 (1981).
- [5] J. Kondo. Physica **84B**. 40 (1976).
- [6] K. Yamada Prog. Theor. Phys. **70**, 73 (1983).
- [7] B. Carmeli and D. Chandler, J. Chem. Phys. **82**, 3400 (1985).
- [8] P. Pfeifer, Ph.D. thesis. Diss. ETH No. 6551, Zurich, 1980.
- [9] N. Kumar and A. M. Jayannavar, Phys. Rev. B **45**, 12014 (1991).
- [10] A. M. Jayannavar, J. Phys. I. France **3**, 1969 (1993).
- [11] H. Spohn and R. Dumcke. J. Stat. Phys. **41**, 389 (1985).
- [12] H. Spohn, Commun. Math. Phys. **123**, 277 (1989).

- [13] P. W. Anderson. Phys. Rev. Lett. **18**, 1049 (1967) , Phys. Rev. **164**, 3 (1967).
- [14] S. Chakravarty and A. J. Leggett, Phys. Rev. Lett. **52** , 5 (1984).
- [15] A. J. Bray and M. A. Moore , Phys. Rev. Lett. **49**, 1545 (1982).
- [16] W. Zwerger. Phys. Rev. Lett. **61**, 1533 (1988).
- [17] T. Tsuzuki, Prog. Theor. Phys. **81**, 770 (1989).
- [18] R. Silbey and R. A. Harris , J. Chem. Phys. **80**, 2615 (1984).
- [19] A. Tanaka and A. Sakurai, Prog. Theor. Phys. **76**, 999 (1986).
- [20] H. Chen and Lu Yu, Phys. Rev. B **45** , 13753 (1992).
- [21] J. P. Sethna. Phys. Rev. B **25** , 5050 (1982).
- [22] H. Chen, Y.Zhang and Xiang Wu, Phys. Rev. B **39**, 546 (1989).
- [23] Y. Zhang,H. Chen and Xiang Wu, J. Phys. C. **2** , 3119 (1990).
- [24] G. Zhang,H. Chen and Xiang Wu, Phys. Rev. B **41**, 11600 (1990).
- [25] B. Dutta. and A. M. Jayannavar , Phys. Rev. B **49**,3604 (1994).

- [26] G. Zhang , H. Chen and X. Wu , ICTP preprint IC/91/223
- [27] G. Zhang , H. Chen and X. Wu. , Phys. Rev. B **43**, 13566 (1991).
- [28] D. F. Walls, Nature (london **306**, 141 (1983))
- [29] A. M. Jayannavar, Solid State Commun. **71**, 689 (1989)
- [30] D. Feinberg,S. Ciuchi and F. dePesqual, Int. J. Mod. Phys. B **4**, 1317 (1990).

FIGURE CAPTIONS

Fig.(1). Plot of reduction factor versus dimensionless dissipation constant α , for a fixed value $\Delta_o/\omega_c = 0.1$. The , - - - - , and ——— lines are obtained from VWF(1), VWF(2) and VWF(3) respectively.

Fig.(2). Phase diagram for the delocalization-localization in $(\Delta_o/\omega_c) - \alpha$ plane.

Fig.(3). The elementary excitation spectrum $(\Omega(k)/\omega_c)$ of the low frequency phonons versus scaled wave vector k/k_c for a fixed value of $\Delta_o = 0.1$, $\alpha = .9599$ and the corresponding value of $Z=0.1005$. The dotted lines(- - -) represent excitation spectrum $\omega_k = |k|$, for uncoupled phonon modes.

Fig.(4). The zero wave vector gap $\Omega(0)/\omega_c$ in the excitation spectrum versus dimensionless dissipation constant α . For a fixed value of $\Delta_o = 0.1$.

Fig.(5). The phonon field displacement spatial correlation function $C(x)$ versus scaled distance $k_c x$ for a fixed value of $\Delta_o = 0.1$, $\alpha = .9599$ and the corresponding $Z=0.1005$.